

2.06 CHEST PAIN / ACUTE CORONARY SYNDROME

Strive for total on-scene time of less than or equal to 15 minutes.

BLS Treatment

- Assess circulation, airway, breathing, and responsiveness.
- Bilateral blood pressures.
- **Oxygen** as indicated.
- Position of comfort.
- **Aspirin.** (Aspirin is in EMT scope of practice)
- NPO. Unless otherwise noted .
- Either list patient medications on PCR or gather medication vials for transport to hospital.

ALS Treatment

Establish a large bore (18G or larger) IV with **Normal Saline** TKO. If possible, establish a second large bore NS lock in the same arm.

12-lead EKG:

- ⊖ Do prior to administration of **Nitroglycerin (NTG)** or pain medication.
- Transmit if EKG interpretation is “STEMI” and notify appropriate STAR center.
- Apply “stand-by” defibrillation pads to all EKG confirmed STEMI patients.

Nitroglycerin: DO NOT administer **NTG** to patients who have taken a phosphodiesterase inhibitor (erectile dysfunction drugs) within the following time frames:

- Sildenafil (Viagra, Revatio) or Vardenafil (Levitra, Staxin) < 24 hours
- Tadalafil (Cialis, Adcirca) < 48 hours
- See “Use 12-Lead EKG to Determine Safety of NTG Administration.”

Persistent chest pain of suspected cardiac origin at any level (scale 1 – 10) shall be treated with **Morphine** or **Fentanyl**. Doses may be started at lower levels than for traumatic or other types of pain treatment.

Ondansetron as needed for nausea.

If hemodynamically unstable, go to Protocol 2.16 Shock.

2.06 CHEST PAIN / ACUTE CORONARY SYNDROME

USE 12-LEAD EKG TO DETERMINE SAFETY OF NTG ADMINISTRATION

- Follow Protocol 7.10 12-Lead EKG.
- Determine presence of ST elevation in leads II, III and AVF. If ST elevation is present, then apply V4R lead.
 - If ST elevation present in V4R, do NOT give **NTG** (to maintain RV filling pressure).
 - If ST elevation in V4R AND clinical signs of shock, including SBP < 90 Hg go to Protocol 2.16 Shock.

Documentation

- “At Patient Side” Time
- VS including bilateral BPs and room air O2 saturation.
- Reassessment of patient symptoms, complaints and vital signs.” At minimum, two sets of vital signs and a reassessment should be done and documented in the PCR after any intervention.
- “O-P-Q-R-S-T symptom assessment:
 - **O** = Onset (Sudden or gradual)
 - **P** = Provoke (What were you doing when the pain started? Does anything make it better or worse?)
 - **Q** = Quality (What does the pain feel like?)
 - **R** = Region/Radiate (Where is the pain? Does it go anywhere else?)
 - **S** = Severity (On a scale of 1-10, 10 being the worst pain you have ever had, how would you rate that pain now? How would you rate that pain at its worst or during exertion/movement?)
 - **T** = Time (When or what time did this start?)
- Aspirin given by EMS. Note if patient self-administered Aspirin or if it was given by someone else (e.g. medical provider).
- EKG findings.
- List patient identifiers on **ALL** transmitted EKGs:
 - Patient Last Name + First Initial
 - Gender
 - Age
 - Ambulance company name and unit number.