

MIDAZOLAM (VERSED®)

THERAPEUTIC EFFECTS	Sedation	
INDICATIONS	<ol style="list-style-type: none"> 1. Adult or pediatric seizures* 2. To facilitate external pacing 3. Sedation following intubation* 4. Excited delirium or severe agitation interfering with necessary patient assessment and/or treatment* 5. Premedication prior to cardioversion or endotracheal intubation if etomidate is not available*† 	
CONTRAINDICATIONS	<ul style="list-style-type: none"> ✓ Hypersensitivity to midazolam ✓ Acute narrow angle glaucoma ✓ Shock 	
PRECAUTIONS/SIDE EFFECTS	May depress respirations or produce over-sedation. Due to a short half-life of the drug, readministration may be necessary.	
ADULT DOSAGE/ROUTE	<p>Indications 1,2,3,5</p> <p>1-2 mg q 3 minutes IV/IM. Up to a max dose of 6 mg</p>	<p>Indication 4</p> <p>2.5 mg q 3-5 minutes IV/IM up to a max dose of 10 mg*</p>
PEDIATRIC DOSAGE/ROUTE	IV/IM/IO .05-0.1 mg/kg q 3 min to a maximum dose of 2 mg†	

*May be given intranasally via Mucosal Atomization Device (MAD) at a dose of 5 mg.

†May be given intranasally via MAD at a dose of 0.2 mg/kg up to a maximum dose of 2 mg.