



Ondansetron/Zofran

CLASS:	Antiemetic
ACTIONS:	Relieves nausea and vomiting
INDICATIONS:	Adult Nausea and Vomiting Adult Pain Management Pediatric Pain Management
CONTRAINDICATIONS:	Known hypersensitivity to the drug
PRECAUTIONS:	Congestive heart failure Bradydysrhythmias Prolonged QT Syndrome Hepatic impairment
SIDE EFFECTS:	Headache Constipation Diarrhea Atrial fibrillation Prolonged QT interval Drowsiness
ADULT DOSAGE:	Nausea/Vomiting: Administer <i>Ondansetron (Zofran) 4 mg IM or IV over 2 min.</i> If vomiting continues, repeat <i>4 mg</i> in 10 minutes. Pain Management: Administer <i>Zofran (Ondansetron) 4 mg IV/IO/IM</i>
PEDIATRIC DOSAGE:	Pain Management: For pediatric patients greater than one year of age and less than 40 kg, administer <i>Zofran (Ondansetron) 0.15 mg/kg slow IV/IO up to a maximum dose of 4 mg.</i> For pediatric patients greater than one year of age and greater than 40 kg, administer <i>Zofran (Ondansetron) 4 mg slow IV/IO.</i>