



Sublimaze/Fentanyl

CLASS:	Synthetic Opiate Analgesic
ACTIONS:	Binds to opioid receptors, producing analgesia and sedation: opioid agonist
INDICATIONS:	Adult Coronary Syndromes Adult Pain Management
CONTRAINDICATIONS:	Systolic BP less than 90 mm/Hg Hypersensitivity to drug
PRECAUTIONS:	Respiratory Depression Bradycardia COPD Elderly and Debilitated Patients MAO Inhibitor Therapy Head Injury
SIDE EFFECTS:	Hypotension Respiratory Depression Nausea and Vomiting Bradycardia Muscle Rigidity
ADULT DOSAGE:	Acute Coronary Syndrome: <i>Fentanyl (Sublimaze) 25 mcg IN, IM or IV/IO over 2 minutes as initial dose; may repeat 25 mcg every 5 minutes titrated to pain relief up to maximum dose of 100 mcg as long as patient systolic blood pressure is greater than 90 mmHg and patient remains conscious</i> Pain Management: <i>Fentanyl (Sublimaze) 25 mcg IN, IM or IV/IO over 2 minutes as initial dose; may repeat 25 mcg every 5 minutes titrated to pain relief up to maximum dose of 100 mcg as long as patient systolic blood pressure is greater than 90 mmHg and patient remains conscious</i>
PEDIATRIC DOSAGE:	Not currently indicated for pediatric use in protocols