




Section 9: Pediatric ACLS Protocols

PEDS ACLS: VENTRICULAR FIBRILLATION (V-FIB) PULSELESS VENTRICULAR TACHYCARDIA





Section 9: Pediatric ACLS Protocols

PEDS ACLS: VENTRICULAR FIBRILLATION (V-FIB) PULSELESS VENTRICULAR TACHYCARDIA-Cont.

PEARLS and KEY POINTS

HISTORY	SIGNS AND SYMPTOMS	DIFFERENTIAL DIAGNOSIS
<ul style="list-style-type: none">• Time of arrest• Medical history• Medications• Possibility of foreign body• Hypothermia	<ul style="list-style-type: none">• Unresponsive• Cardiac arrest	<ul style="list-style-type: none">• Respiratory failure• Foreign body• Secretions• Infection (croup, epiglottitis)• Hypovolemia (dehydration)• Congenital heart disease• Trauma• Tension pneumothorax• Hypothermia• Toxin or medication• Hypoglycemia• Acidosis

Do Not Confuse Epinephrine 1:1000 IM dose and 1:10,000 IVP dose

- **Exam: Mental Status**
- Monophasic and Biphasic waveform defibrillators should use the same energy levels noted.
- In order to be successful in pediatric arrests, a cause must be identified and corrected.
- Airway is the most important intervention. This should be accomplished immediately. Patient survival is often dependent on airway management success.
- If the patient converts to another rhythm, follow the appropriate protocol and treat accordingly.
- If the patient converts back to ventricular fibrillation or pulseless ventricular tachycardia, defibrillate at the previously used setting.
- Defibrillation is the definitive therapy for ventricular fibrillation and pulseless ventricular tachycardia.
- Defibrillate 30 - 60 seconds after each medication administration.
- The proper administration sequence is shock, drug, shock, and drug.