

Section 13: Appendix 1: Medications

NITROGLYCERINE (NITRO-STAT)

Pregnancy Category - C

Е	EMT	Е
Α	AEMT	Α
P	PARAMEDIC	P

ACTIONS	1. Vasodilatation
	Coronary artery dilation
	Decreases myocardial oxygen demand
	Decreases vascular resistance
INDICATIONS	Suspected ischemic chest pain / AMI
	Hypertensive emergency with signs and symptoms of ACS
	Pulmonary edema
CONTRAINDICATIONS	1. Hypotension
	Known hypersensitivity
	3. Use of Viagra or similar erectile dysfunction medications within
	48 hours
PRECAUTIONS	1. Use caution in patients with inferior wall MI (Elevation in leads II,
	III, AVF)
	Avoid use in patients with increased intracranial pressure or
	glaucoma
	If the patient becomes hypotensive after nitroglycerine
	administration, then place the patient in a semi-reclined position
	with legs elevated and give IV normal saline bolus
SIDE EFFECTS	1. Hypotension
	Throbbing headache
	Lightheadedness / dizziness
	4. Syncope
SUPPLIED	0.4 mg SL tablet
ADULT DOSAGE	Cardiac Chest Discomfort / AMI:
	0.4 mg SL (may be repeated up to 3 doses total)
	EMT CONTACT MEDICAL CONTROL
	Pulmonary Edema / CHF:
	0.4 mg SL (may be repeated up to 3 doses total)
	Fourthernal Fourier Body
	Esophageal Foreign Body:
DEDIATRIC DOCACE	0.4 mg SL
PEDIATRIC DOSAGE	Not recommended in prehospital setting
KEY POINTS	 May repeat up to 3 doses if B/P systolic > 110 with IV or 120 without
	IV
	Assure that patient does not chew or swallow tablets
PROTOCOL USE	Acute Coronary Symptoms
	Adult Esophageal Foreign Body Obstruction