



Section 13: Appendix 1: Medications

BRILINTA (TICAGRELOR)

Pregnancy Category - C

P PARAMEDIC P

ACTIONS	<ol style="list-style-type: none">1. Platelet P2Y12 inhibitor that reduces the rate of thrombotic events in ACS and STEMI patients2. Reduces reinfarction in STEMI or Stroke patients
INDICATIONS	<ol style="list-style-type: none">1. STEMI Patients-ST Elevation myocardial infarction
CONTRAINDICATIONS	<ol style="list-style-type: none">1. History of Stroke2. GI Bleed3. Trauma4. Intracranial hemorrhage5. Age 80 or older6. Patient takes Coumadin, Pradaxa, Eliquis, xarelto or savaysa
PRECAUTIONS	<ol style="list-style-type: none">1. Active pathologic bleeding may occur (peptic ulcer)2. Use with caution in patients with liver disease3. Use with caution following major surgery. Can cause fatal bleeding
SUPPLIED	90 mg tablet
SIDE EFFECTS	<ol style="list-style-type: none">1. Hemorrhage2. Dyspnea3. Headache4. Dizziness
ADULT DOSAGE	180 mg (two 90 mg tablets)
PEDIATRIC DOSAGE	Not Indicated in the pre-hospital setting
KEY POINTS	<ul style="list-style-type: none">• Dyspnea can occur (14%) but is self limiting
PROTOCOL USE	<ul style="list-style-type: none">• Adult ACS Protocol