



Yolo County Emergency Medical Services Agency

Protocols

Revised Date: September 1, 2018

SEIZURE	
Adult	Pediatric
BLS	
Assess vital signs O ₂ , titrate SpO ₂ to ≥ 94% Assist ventilations as needed Temperature Cooling measures if febrile	
BLS Local Scope	
Blood Glucose Check	
ALS	
Cardiac Monitor, Waveform EtCO ₂ , Vascular Access	
<p>Midazolam 2mg SIVP</p> <ul style="list-style-type: none"> May repeat at 1 – 2 mg increments every 3 minutes Total max dose 6 mg <p style="text-align: center;"><u>Or</u></p> <p>Midazolam 5 mg IM</p> <ul style="list-style-type: none"> No repeat <p style="text-align: center;"><u>Or</u></p> <p>Midazolam 10 mg IN</p> <ul style="list-style-type: none"> ½ dose each nare No repeat 	<p>Midazolam 0.1 mg/kg SIVP</p> <ul style="list-style-type: none"> May repeat x 2 every 5 - 10 minutes Max single dose 2 mg Total max dose 4 mg <p style="text-align: center;"><i>*Use a 1 mL syringe for Midazolam administration in pediatric patients</i></p> <p style="text-align: center;"><u>Or</u></p> <p>Midazolam 0.1 mg/kg IM</p> <ul style="list-style-type: none"> Max dose 4 mg No repeat <p style="text-align: center;"><u>Or</u></p> <p>Midazolam 0.2 mg/kg IN</p> <ul style="list-style-type: none"> ½ dose each nare Max dose 5mg No repeat
Direction	
<ul style="list-style-type: none"> Contact Receiving ED Physician for additional treatment 	